

To ensure conservation of Minnesota's dragonflies and damselflies through research, education, and outdoor recreation. An effort to determine the ranges and distributions of the dragonflies and damselflies of Minnesota



2006-2012



Historical Collections




Targeted Surveys

Survey Workshops With Volunteers




"A Minnesota-shaped Hole"



• Minnesota range of the Skimming Bluet.

Minnesota Biomes



Methods Part 1


- Determine the historical record by cataloguing the known collections at universities, museums, and other sites.
- UMSP
- Science Museum of MN
- Warner Nature Center
- Carleton College
- Gustavus Adolphus
- St Olaf College
- Winona State University
- UM Morris



Methods Part 2

Targeted Surveys





Methods Part 3

- Citizen Science
- In seven years we held 29 one-day survey workshops and 7 weekend-long Minnesota Dragonfly Gatherings
- An average of 17 participants per workshop and an average of 45 participants for the weekend events in the last two years.

Results

- 23 Odonata species added to the Minnesota state list.
- 2 species removed from the state list.
- 2580 specimens collected by the MOSP
- During the 7 years of the MOSP, 1881 new county records were generated in Minnesota.
- Current total Minnesota Odonata list sits at 148 species

Post-2013 Damselflies

10 species of Damselflies have been added to the Minnesota List since 2006

Red Damsel

Amphiagrion sp.

- Seep-fed marshes and spring-fed streams
- One museum
 specimen from 1935
- Rediscovered in 2008

Kurt Mead

Great Spreadwing

Archilestes grandis

- Small, wooded streams
- First photographed in 2010
- Found in two counties in 2011



Violet Dancer

Argia fumipennes violaceae

Kurt Mead

- Museum specimen from 1935
- Rediscovered in 2011
- Widespread and common

Springwater Dancer Argia plana

- Common through the Great Plains, SW to Mexico
- Discovered at Blue Mounds State Park in 2011






- Museum specimen from 1965
- Rediscovered in 2006

River Bluet Distribution in the Midwest

(map from odonatacentral.com, June 22, 2018)



Azure Bluet Enallagma aspersum

- Discovered in 2009
- Requires fishless waters



Skimming Bluet Enallagma geminatum

- Discovered in 2006
- Now widespread throughout Minnesota



Vernal Bluet Enallagma vernale

Discovered in 2013

Possibly widespread,
 but easily confused with
 other species

Citrine Forktail

Ischnura hastata

- Museum specimen from 1968
- Rediscovered in 2010
- Probably irruptive in Minnesota



Southern Spreadwing Lestes australis



- Discovered in 2011, widespread
- Newly recognized species (was subspecies of a common species)

Post-2013 Dragonflies

13 Dragonfly species have been added to the Minnesota List since 2006

Harlequin Darner Gomphaeschna furcillata



• One specimen from Minnesota in 2012.

Spatterdock Darner Rhionaeschna mutata



- One isolated population in Washington County, breeding in one pair of ponds
- Require fishless waters

Blue-eyed Darner Rhionaeschna multicolor

- Common in the western US
- Only found in eastern MN
- Why not in prairies?



Cyrano Darner Nasiaeschna pentacantha

Scattered records throughout the northern half of Minnesota



Sioux Snaketail Ophiogomphus smithi




- First described in 2004
- Dependant on clean, sandy bottomed rivers
- Discovered in 2011 in upper reaches of the Zumbro River in SE MN
- Specimens from the 1990's uncovered in museum collections from Chisago County

Jade Clubtail Arigomphus submedianus



Two photo records from the Twin Cities

Royal River Cruiser Macromia taeniolata



* Data from Odonata Central



In Minnesota, found only from the Mississippi River in SE counties.


Quebec Emerald Somatochlora brevicincta



 Breed in patterned fens, a very rare habitat type in MN.

Quebec Emerald Distribution

(map from odonatacentral.com)



Ringed Emerald Somatochlora albicincta

- One adult specimen from Lake County
- Efforts targeting the adult and nymph have turned up nothing. We have the habitat...

Ebony Boghaunter Williamsoni fletcheri



 Breed in tiny bog pools and emerge very early in the spring. Slaty Skimmer Libellula incesta

Great Blue Skimmer

Striped Saddlebags Tramea calverti

• All are likely strays to Minnesota

What species are still hiding in Minnesota?



Or should we ask...

What species are still hiding in Ohio?





To ensure conservation of Minnesota's dragonflies and damselflies through research, education, and outdoor recreation.